

**Qualitative and Nutritional Differences in Processing Tomatoes Grown Under
Commercial Organic and Conventional Production Systems**

Diane M. Barrett^{1*}, Craig Weakley², Jerome V. Diaz¹ and Mitchell Watnik³
¹Department of Food Science & Technology, University of California - Davis
²Small Planet Foods, Inc.

³Department of Statistics, California State University, East Bay

*Corresponding author.

1 **ABSTRACT**
2
3

4 Organically grown products experienced a doubling in percent penetration of organic sales
5 into retail markets during the period from 1997-2003, however there is still a debate over the
6 perceived quality advantage of organically grown fruits and vegetables. In a study focusing on
7 commercial production of processing tomatoes, samples were analyzed from four growers with
8 matched organic and conventional fields. For the four growers studied, individual ANOVA results
9 indicated that tomato juice prepared from organically produced tomatoes on some farms was
10 significantly higher in soluble solids (°Brix), higher in consistency and titratable acidity, but lower
11 in red color, ascorbic acid and total phenolics content in the microwaved juice. Results were
12 significantly different between specific growers, and this may be attributed to differences in soil
13 type and soil nutrients, tomato cultivar, environmental conditions, or other production-related
14 factors. Higher levels of soluble solids, titratable acidity and consistency are desirable for the
15 production of tomato paste, in that tomatoes with these attributes may be more flavorful and require
16 less thermal treatment. This has potential to result both in cost savings from less energy required in
17 paste manufacture, and potentially a higher quality product due to less thermal degradation of color,
18 flavor and nutrients. Future work may involve a larger number of commercial growers, and
19 correlation to controlled university research plots.
20

21 **Keywords:** organic, conventional, processing, tomatoes, nutrition, quality
22

1 INTRODUCTION

2
3
4 In a recent study on factors affecting food choice with respect to fruit and vegetable intake
5 (Pollard and others 2002) it was found that personal ideologies resulted in consumers buying an
6 increasing percentage of organic products, despite rising costs. Of consumers who buy organic
7 produce, a survey commissioned by the Soil Association and Baby Organix (The Soil Association
8 1999) found that 43% of people bought organic because of preferred taste, 28% because it is
9 environmentally friendly and 24% because it is animal welfare friendly. In addition, many
10 consumers perceive that organically grown products are more healthy than conventional, although
11 the research on this aspect is not conclusive.

12 There is scientific evidence that organically grown crops contain higher mineral and vitamin
13 content (Worthington 1998; Diver 2000), higher antioxidant content (Woese and others 1997;
14 Weibel and others 2000; Heaton 2001; Asami and others 2003; Chassy and others 2006), and better
15 flavor (Weibel and others 2000; Reganold and others 2001) than crops produced using conventional
16 production systems. In addition, there is concern by some authors that the antioxidant content of
17 foods grown using conventional production systems is lower than optimal for human health (Woese
18 and others 1997).

19 Other studies have resulted in conclusions that differences in quality and nutritive value of
20 organically and conventional grown foods are inconsistent or unsubstantiated (Woese and others
21 1997; Worthington 1998; Diver 2000; Bourn and Prescott, 2002). In recent reviews of quality
22 comparison studies between organic and conventional foods, Woese and others (1997),
23 Worthington (1998), Brandt and Molgaard (2001), Heaton (2001) and Bourn and Prescott (2002)
24 suggest that interpretation of the results of these investigations is difficult because of
25 methodological differences related to cultivar selection, growing conditions, and sampling and
26 analytical methods. In a recently published Scientific Status Summary by the Institute of Food
27 Technologists (Winter 2006), the author concludes that it is premature to say that either organic or
28 conventional food systems is superior with respect to safety or nutritional composition.

29 Although the quality and potential nutritional benefits of organically grown foods are still
30 being studied, the Organic Trade Association has reported that recent growth in sales of organic
31 products has been 20% per year (Organic Trade Association 2004). The most common point of
32 entry for consumers new to organic products are fruits and vegetables. The Economic Research
33 Service reports that there has been a doubling in the percent penetration of organic sales into retail
34 markets in the most recently documented six year period, e.g. from 2% in 1997 to 4% in 2003
35 (Economic Research Service 2005a).

36 According to the California Department of Food and Agriculture (Klonsky and Richter, in
37 press), tomatoes are sixth in the top 10 organically grown commodities produced in the state, with
38 the combination of fresh and processing tomato varieties representing \$11 million in sales in 2003
39 (Economic Research Service 2005b). Ninety-three percent of the organic tomatoes sold in the U.S.
40 in 2003 originated from California farms (Economic Research Service 2005a). Acreage of
41 processing tomatoes in California has increased from 2,232 to 4,108 in the period from 1998 to
42 2003 (Klonsky and Richter, in press) and the number of organic growers has risen in the same
43 period from 33 to 41. Farm level sales of organically grown processing tomatoes have gone from
44 \$4.6 million in 1998 to \$5.3 million in 2003 (Economic Research Service 2005b).

45 The objective of this study was to compare the quality and nutritional value of processing
46 tomatoes grown on matched commercial grower fields in California. Several tomato fruit quality
47 components, including °Brix, pH, titratable acidity, color, ascorbic acid content, total phenolic
48 content, consistency, and sensory quality were compared in processing tomatoes grown by
49 conventional production systems and certified organic production systems. Commercial scale
50 processing tomato growers experienced in growing organic tomatoes were collaborators on these

1 trials, and soil samples were taken both prior to planting in order to substantiate that soil types on
2 organic and conventional fields on the same farm were similar. Soil samples were also taken after
3 fruit set in order to determine the correlation between soil characteristics and fruit quality.
4
5

6 **MATERIALS AND METHODS**

7
8

9 Grower Field Statistics and Selection of Sites. In 2003, tomatoes were grown on four pairs of
10 matched fields in the California Central Valley. Each field pair consisted of matched organic and
11 conventional fields. In order to assure that soil type, topography, and other dominant soil
12 characteristics were the same for each field pair, research sites were selected using United States
13 Department of Agriculture-Natural Resources Conservation Service (USDA-NRCS) soil scientist
14 evaluations prior to planting. The soil profile in each field was examined using soil sample cores to
15 a depth of at least 102 cm (Table 1). This allowed research sites to be selected so that soil type,
16 topography, and other soil characteristics would not likely be significant factors in the comparison
17 of tomato fruit quality components between field pairs. Each matched field pair of organic and
18 conventional tomatoes had the same grower, same soil type, same tomato cultivar, same planting
19 date, same irrigation method, and a very similar harvest date (Table 2). Three growers (Harris,
20 O'Neill and Terranova) from Fresno county, and one grower (Rominger) from Yolo county,
21 representing major processing tomato regions in California, participated in this study.
22

23 Production Inputs. Design of this study attempted to control for grower skill, soil type, tomato
24 cultivar, planting date, irrigation method, and harvest date within each field pair, the production
25 systems were substantially different between farms (Table 3). All organic fields in this study were
26 farmed in accordance with the USDA National Organic Standards. The organic crop production
27 methods included: 1) soil building with compost, manure, and/or cover cropping; 2) weed control
28 with mechanical cultivation and hand hoeing; 3) plant disease control with copper and sulfur; and
29 4) insect pest management with biological control methods. The conventional crop production
30 methods included: 1) fertilization with synthetic chemical fertilizers (NPK); 2) weed control with
31 synthetic chemical herbicides; 3) plant disease control with copper, sulfur, and synthetic chemical
32 fungicides; and 4) insect pest management with synthetic chemical insecticides.
33

34 Research Plot Designation. Each research area was eleven beds wide (1.52 to 1.68 meter beds) by
35 91.5 meters long; approximately 0.2 hectares in size. Research areas were oriented in the same
36 north-south-east-west direction in each field pair. Within each research area, 88 plots were
37 established with each plot being one bed wide by 0.12 meters long. Eight plots within each field's
38 research area were randomly selected for data collection. The selected plots were matched for each
39 field pair.
40

1 Soil Sampling. Prior to choosing the organic and conventional fields to be used at each grower
2 location, soil samples were taken from four randomly selected research plots within each field's
3 research area to determine whether organic and conventional fields were properly matched by soil
4 type. Tomatoes were planted in closely matched fields, and after fruit set, additional soil samples
5 were taken for comparison to and correlation with fruit quality. In each plot, twelve sub-samples
6 were taken with an Oakfield soil sampling tube (1.9 cm diameter) from a depth of 0 to 10.2 cm.
7 Soil sub-samples were composited, air dried, and analyzed by the Division of Natural Resources
8 (DANR) Analytical Laboratory at the University of California - Davis. Methods may be found on-
9 line at <http://danranlab.ucanr.org>.

10 Soil salinity and alkalinity measurements included pH, cation exchange capacity (CEC),
11 estimated soluble salts, sodium absorption ratio (SAR), exchangeable sodium percentage (ESR). A
12 saturated paste extract was prepared and the following were also measured: HCO₃, Ca, Mg, Na, Cl.
13 Soil fertility measurements included soil nitrate and extractable ammonium, total Kjeldhal nitrogen
14 (TKN) and extractable potassium using the Olsen method. The following micronutrients were
15 extracted with diethylenetriaminepentaacetic acid: Zn, Mn, Fe, Cu. In addition, determination of
16 exchangeable minerals (X-K, X-Ca, X-Mg, X-Na) were carried out. The following physico-
17 chemical characteristics were evaluated: organic matter, particle size analysis (sand/silt/clay).
18

19 Tomato Sample Harvesting. At harvest, 4.54 kg of red ripe fruit were hand harvested from each of
20 the eight research plots in each field (Table 2) and delivered the same day to the University of
21 California, Davis, Food Science and Technology laboratory for analysis of fruit quality
22 components. In addition, approximately 5.44 kg of red ripe tomatoes were collected at a later date
23 from each of four randomly selected research plots in each field and delivered for sensory analysis.
24 Sensory samples were collected on the same date for each field pair.
25

26 **Tomato Quality Analysis**

27
28 Moisture Content. Moisture content in the raw tomatoes was analyzed according to the Association
29 of Official Analytical Chemists (AOAC) vacuum oven method (AOAC International 2000) on
30 organic and conventionally grown tomatoes. Triplicate moisture measurements were made on
31 approximately 3g samples from each of the 4.54 kg samples harvested.
32

33 Microwave Hot Break Process. A microwave hot break method developed in the Department of
34 Food Science & Technology at the University of California – Davis (Leonard and others 1980) was
35 used as a rapid means of simulating an industrial hot break, or thermal step that is utilized to
36 inactivate enzymes. Enzyme inactivation serves to stabilize the tomato juice to a certain degree,
37 and prevent enzyme-catalyzed changes in viscosity, color, flavor and nutrient content. Predictive
38 equations were previously developed by Leonard and others (1980) to correlate measurements on
39 microwaved tomato juice to catsup yield. Catsup yield is calculated using the following equations
40 (Leonard and others 1980):

41 $\text{Paste yield} = 908 \text{ kg} * \text{tomato juice } ^\circ\text{Brix} / 28 ^\circ\text{Brix}$

42 $\% \text{ tomato solids} = 7.388 + 1.015(\text{paste Bostwick}) - 0.0138(\text{paste Bostwick})$

43 $\text{Catsup yield} = 908 \text{ kg} * \text{tomato juice } ^\circ\text{Brix} / \% \text{ tomato solids}$
44

45 Tomatoes were washed, towel dried and sorted for defects, then cut in half for determination
46 of quality factors on the microwave break juice. One half of each fruit was placed in a Pyrex dish
47 to achieve a net weight of approximately 1300 grams. The dish was immediately weighed, covered
48 and microwaved in a commercial (1400 watt) microwave oven for 6 min at 100% power, followed
49 by 6 min at 50% power. After cooking, the dish was placed in ice water to cool. Cooled samples

1 were re-weighed, and water was added to compensate for evaporative losses during cooking. Seeds
2 and skins were extracted using a lab pulper with a 0.033 inch screen.

3
4 Titratable acidity, pH, °Brix, Bostwick and Color Measurements. Following the microwave
5 treatment, pulped juice samples were evaluated for titratable acidity using titration with NaOH
6 (AOAC International 2000). The remaining juice was deaerated and the temperature adjusted to 25
7 $\pm 0.2^\circ\text{C}$, then used for determination of pH and °Brix (soluble sugars). Independent duplicate
8 Bostwick consistency readings were obtained on each sample (Barrett and Anthon 2001). Readings
9 reported are the distance (cm) that a volume of juice of fixed dimension flowed in a trough in 30
10 sec. A smaller reading corresponds to less flow, or product of higher consistency.

11 Color (HunterLab, Reston, VA) values were also measured on microwaved deaerated juice.
12 Instrumental color measurements were made on juice samples placed in glass sample cups. L value
13 (white to black or light to dark), a (green to red) and b (yellow to blue) measurements were taken
14 with a Hunter colorimeter. The colorimeter was calibrated with a white tile and a standard tile of a
15 color similar to that of the sample. L, a and b values were determined by averaging the results of
16 three independent readings per sample. From the L, a, b values, USDA tomato scores were
17 calculated. In addition, overall color was measured on microwaved deaerated juice using a Light
18 Emitting Device (LED), a standard colorimeter used by the California tomato (Valero and others
19 2003).

20
21 Lycopene. A modification (Barrett and Anthon 2001) of the method published by the AOAC (2000)
22 was used for lycopene analysis. First 100 μl of microwaved tomato juice was pipetted into a screw
23 cap tube using a 100 μl Drummond micropipettor. Then 7.0 ml of 4:3 (v/v) ethanol:hexane was
24 added, the tube was capped, vortexed, then incubated, out of bright light, with occasional vortexing.
25 After 1 hr 1.0 ml water was added to each sample and then shaken briefly. Samples were allowed
26 to stand 10 minutes to afford phase separation and dissipation of air bubbles. A sample of the
27 hexane layer was read at Abs 503 versus hexane in the spectrophotometer (Shimadzu, Japan).
28 Lycopene levels in the hexane extracts were then calculated according to:

$$29 \quad \mu\text{g lycopene/g fresh wt.} = (A_{503} \times 537 \times 2.7)/(0.1 \times 172) = A_{503} \times 84.3$$

30 where 537 g/mole is the molecular weight of lycopene, 2.7 ml is the volume of the hexane layer, 0.1
31 g is the weight of sample added, and 172 mM^{-1} is the extinction coefficient for lycopene in hexane.
32 Duplicate samples were analyzed.

33
34 Ascorbic (reduced) and Dehydroascorbic (oxidized) Acid. Raw and microwaved tomato juice
35 samples were analyzed for ascorbic acid, dehydroascorbic acid and total phenolics. Ascorbic acid
36 was determined using a spectrophotometric method (Latapi and Barrett 2006). One gram of sample
37 was homogenized with distilled water using a Polytron (Brinkmann Instruments Inc., Model
38 PCU11, Westbury, NY) until a thick paste was obtained. The paste was centrifuged and the
39 supernatant removed for analysis. In a 3 ml cuvette, 2.5 ml of 0.1M sodium phosphate, pH 6.5, 0.1
40 ml of sample, 0.4 ml of water, and 0.5 ml of 1.0 mg/ml horseradish peroxidase (Sigma Type II)
41 were mixed. The sample was read at Abs 265 in a spectrophotometer (Shimadzu, Japan) to
42 determine total ascorbic acid, 50 mM hydrogen peroxide was added and the sample was read again
43 after it reached a stable absorbance value at 265 nm to determine oxidized or dehydroascorbic acid.
44 Ascorbic acid content was expressed as mg/g dry weight. Measurements were performed in
45 triplicate.

46
47 Total Phenolics. Total phenolics concentrations were measured on raw and microwaved tomato
48 juice using the Folin-Ciocalteu assay. Five ml acetone, 0.5 ml sample, and 1.0 ml Folin-Ciocalteu
49 reagent were added to a 25 ml volumetric flask. The contents were mixed and allowed to stand for
50 5 to 8 min at room temperature. Ten ml of 7% sodium carbonate solution was added, followed by

1 the addition of nanopure water filled to volume. Solutions were mixed and allowed to stand at
2 room temperature for 2 h. Sample aliquots were filtered through a Whatman 0.45 um
3 polytetrafluoroethylene filter prior to the determination of total phenols concentration using a
4 spectrophotometer (Shimadzu, Japan) monitoring Abs 750 nm. Total phenolics content was
5 standardized against gallic acid and expressed as gallic acid equivalents (GAE). The linearity range
6 for this assay was determined as 0.5 to 5.0 mg/ μ L GAE ($R^2 = 0.9990$) giving an absorbance range
7 of 0.050 to 0.555 AU.

8
9 Diced Tomato Texture. Approximately 10 tomatoes were diced and mixed, and firmness was
10 analyzed (Garcia and Barrett 2006) in triplicate 200 gram samples using a Kramer shear cell and a
11 TA.XT2 Texture Analyzer (Texture Technologies, Scarsdale, NY). Average values were
12 calculated.

13
14 Sensory Evaluation. Fresh, uncooked, diced tomatoes were analyzed for sensory quality using
15 consumer preference tests. Samples were presented at random to over one hundred untrained
16 panelists. Samples were evaluated for degree of liking on the 9 point Hedonic scale where: 1 =
17 dislike extremely, 5 = neither like nor dislike, and 9 = like extremely. Panelists were asked to
18 comment on liking of color, texture, flavor and overall quality. Data was analyzed by analysis of
19 variance and the Fisher's Least Significance Difference Test was used to compare differences
20 among means at the 0.05 level (Microsoft Excel (2000) for Windows computer software).

21
22 Evaluation of Tomato Peelability and Product Yield. Six batches of 20 clean, sorted tomatoes were
23 weighed for each lot of fruit and peeled by exposure to live steam (30 psig) for 45 s in a pilot scale
24 Odenberg (Odenberg, West Sacramento, CA) steam peeler. Following steam exposure, tomatoes
25 were passed over pilot scale disc and pinch rollers. Peelability was classified using a subjective
26 rating system (Barrett 2001) and those with little or no peel were designated as "peeled". Peeled
27 fruits were weighed and % whole peel yield was determined. Values of % peeled and % whole peel
28 yield for the six replicate processing batches were averaged.

30 **Statistical Analysis**

31
32 Data were first analyzed using Multivariate Analysis of Variance (MANOVA) models
33 within SAS (SAS Institute, v. 9.1, 2002, Cary, NC) "proc glm". MANOVA models differ from
34 regular ANOVA models in that there is more than one outcome. In MANOVA models there is less
35 probability of false claims of statistical significance, and the models are consistent for all outcomes
36 (Johnson and Wichern 1998). Multivariate F values (Wilk's Lamda, Pillai's Trace, Hotelling
37 Lawley Trace and Roy's Greatest Root) were obtained based on a comparison of the error
38 variance/covariance matrix and the effect variance/covariance matrix. Growers were treated as a
39 "fixed" effect, that is, the results are true for the specific growers that were studied and cannot be
40 generalized to the general population of conventional or organic tomato growers. In order to
41 generalize results to the general population, the growers would need to be treated as a random
42 effect. In this study, the use of growers as a random effect would have provided low statistical
43 power for the effects of interest.

44 After doing the overall MANOVA to see general trends and eliminating insignificant terms,
45 we looked at individual quality parameters using individual or "protected" ANOVAs to determine
46 whether they contributed to the overall significant difference. Means comparisons were performed
47 using Least Squares (LS) Means tests (SAS, 2002). This allowed us to compare organic and
48 conventional production systems at individual grower locations. We used unadjusted comparisons
49 to maintain statistical power.

1 Twenty-two different soil characteristics were analyzed. We provide comparisons between
2 growers for each of these characteristics. It was also of interest to determine whether the soil
3 characteristics could account for any observed differences in tomato quality attributes. However,
4 because soil samples were taken from only four field locations, it was only possible to use the same
5 four (rather than all eight) location-specific samples taken for quality analysis. Principal
6 component analysis (PCA) was used to reduce the number of soil variables to those that were best
7 able to discriminate the four growers and those that discriminated conventional and organic
8 production practices (Johnson and Wichern 1998). When quantitative explanatory variables are
9 added to a MANOVA model, the resulting model is referred to as Multivariate Analysis of
10 Covariance (MANCOVA).

11 12 13 **RESULTS AND DISCUSSION**

14
15 The four grower fields utilized in this study differed in soil type and texture (Table 1), with
16 sandy loam soil in the Harris and Terranova farms and clay soil in the O'Neill and Rominger farms.
17 There were also differences in the tomato cultivar planted, planting type and irrigation method
18 (Table 2) between growers. The cultivar was selected by each specific grower, and in this study a
19 different cultivar was planted by each of the grower participants. O'Neill direct seeded while the
20 other three growers planted transplants. In addition, there were differences in the irrigation
21 methods utilized, with furrow irrigation predominating, but drip irrigation being used at Harris
22 Farms. In terms of the production systems utilized, there were greater similarities in the production
23 systems (whether organic or conventional) at a particular grower location than within the same
24 production system (either organic or conventional) at different grower sites. This illustrates the
25 challenge with conducting research on commercial farms. While scientists would like to evaluate
26 successful production systems in a realistic environment, it is not practical to control production
27 variables and one must acquire whatever information is possible within the constraints of grower
28 prerogatives. Because there were only 4 growers included in this study, it was not possible to make
29 global statements concerning the differences between quality of organic and conventionally grown
30 tomatoes. Rather, the results discussed below will apply only to the four specific growers involved
31 in this study.

32 Table 3 illustrates that even within commercial organic or conventional production systems,
33 there were differences from grower to grower. For the most part, conventional production systems
34 utilized urea ammonium nitrate as a fertilizer, but there were differences in pest control materials
35 and rate of application. The organic soil fertility materials added differed more, with most growers
36 using compost and chicken manure, one grower using turkey manure, and another grower a
37 combination of cow and chicken manure. There were also differences in grower yield (Table 2),
38 but it was not possible based on the data collected to conclude whether organic or conventional
39 production systems yielded more kilograms per hectare. Although we attempted to harvest
40 tomatoes at approximately 90-95% field maturity, there were some instances, in particular with
41 Terranova Ranch and O'Neill Farming Co. (Table 2), where we were required to hand harvest
42 before the field was 90-95% mature because growers had made a decision to machine harvest.

43 Most of the quality analyses carried out on the tomato samples were done following a
44 microwave hot break treatment. This treatment simulates a commercial tomato paste hot break
45 treatment, which targets the inactivation of endogenous enzymes that may affect quality, in
46 particular the product consistency. While heating the tomato juice inactivates polygalacturonase,
47 pectin methylesterase and other quality-related enzymes, it can result in degradation of color and
48 nutrients. For this reason, analysis of ascorbic acid, dehydroascorbic acid and total phenolics was
49 carried out on both raw and microwaved tomato juice.

1 The overall focus of this project was to determine if qualitative and nutritional differences
2 existed between organically grown and conventionally grown processing tomatoes. Analysis of the
3 MANOVA F values (Table 4) indicated that, for all parameters determined, both grower and
4 production system (organic and conventional) were significant, as well as the interaction term,
5 production system x grower. This means that there were differences between fruit produced by
6 organic and conventional production systems, as well as between the four grower participants.
7 Therefore it was not possible to pool growers when evaluating the effect of production system nor
8 was it possible to conclude that organically grown (or vice-versa, conventional grown) tomatoes
9 from one grower would have the same quality attributes as those grown by another grower.

10 Individual ANOVAs carried out using the general linear model on each quality variable
11 (average of eight field locations for each grower and production system) determined the Type III
12 sum of squares (SAS, 2002) values for grower, production system and the interaction term
13 production system x grower (data not shown). Because some of the data did not conform to the
14 statistical assumption of normality of the errors, statistical transformations (log, rank, squared
15 values) were used and are indicated in parentheses. We used 0.05 as our level of statistical
16 significance. It must be stressed that the conclusions from these individual ANOVAs apply to the
17 organic and conventional production systems in the four growers that participated in this study, and
18 may not be extrapolated to the general population of growers.

19 Tables 5, 6 and 7 summarize the quality analyses made on organic and conventionally
20 grown fruit, and the p values for grower, production system and the production system x grower
21 interaction. This interaction was significant ($p < 0.05$) for a number of attributes including °Brix,
22 pH, titratable acidity, catsup yield, a and b color values, USDA color, dehydroascorbic acid in raw
23 tomato juice, ascorbic and dehydroascorbic acid in microwaved juice, lycopene, and phenolics
24 levels both in raw and microwaved juice. Other as of yet unknown factors must have caused these
25 differences.

26 On examination of the grower statistics at the bottom of Tables 5, 6 and 7, it is clear that
27 there were significant differences ($p < 0.05$) between growers in °Brix, pH, titratable acidity, catsup
28 yield, LED color, L, a and b color values, USDA color, ascorbic and dehydroascorbic acid in the
29 raw tomato juice, ascorbic acid and total phenolics in the microwaved tomato juice and lycopene.
30 Grower-specific differences that may affect tomato quality include microclimate, soil type, soil
31 nutrients, irrigation method and moisture-holding capacity of the soil, number of years in
32 cultivation, tomato cultivar, or other unknown factors. In the cases noted above, this study was
33 unable to determine why some quality attributes differed between growers.

34 There were significant differences ($p < 0.05$) between organic and conventional production
35 systems in °Brix, titratable acidity (expressed as % citric acid), Bostwick consistency, catsup yield,
36 LED color, Hunter b value, USDA color, dehydroascorbic acid in raw tomato juice, and ascorbic
37 acid and total phenolics in the microwaved juice. This means that for these quality attributes there
38 was a significant difference between tomatoes grown organically and those grown conventionally.
39 The results from this study of four growers indicated that organically grown tomatoes on some
40 farms were significantly higher in °Brix (positive), higher in titratable acidity (positive) and lower in
41 Bostwick (positive), higher in catsup yield (positive), lower in LED color (negative), lower in b
42 value (positive), lower in ascorbic acid (negative) and phenolics (negative) in the microwaved juice.

43 Four different tomato cultivars were utilized in this study, a different cultivar at each grower
44 location. While this is a good test of the effect of production system on a range of processing
45 tomato varieties, it resulted in a confounding of cultivar and grower, and it may have been better
46 from a statistical point of view to use the same cultivar at all grower locations. Woese and others
47 (1997) stated that, in the case of apples grown under conventional and organic production systems,
48 the differences between cultivars had a far greater influence on the composition of the apples than
49 the different cultivation forms.

50

Grower Specific Differences in Physico-Chemical Properties of Conventionally and Organically Grown Tomatoes

Where overall F values were significant, pairwise comparisons between the organic and conventional production system at each grower location were made using Least Square Means. These results are presented in Tables 5, 6 and 7. While general trends for the four grower participants may be obtained from inspection of the ANOVAs for individual quality attributes, it is necessary to use LSD values to make specific comparisons between organically and conventionally grown tomatoes at each individual grower location.

There were no significant differences in sensory color, texture, flavor or overall quality, diced tomato texture or tomato peelability and product yield when organically grown and conventionally grown tomatoes were compared. Diced tomato texture values (Table 5) ranged from 155 to 300 N, with a mean of 210 N, in the organic tomatoes, and from 188 to 412 N, with a mean of 260 N, in the conventionally produced tomatoes. The conventionally produced tomatoes were somewhat firmer at each farm site, and in the cases of Terranova and Rominger, these were statistically significant differences.

In comparing the peelability and whole peeled product yield, there were no significant differences between the two production systems (data not shown). Percentage of tomatoes peeled ranged from 20 to 95 percent, but replicate values from the same grower and production method were similar. Tomatoes grown under either production method at the Terranova farm had the highest percentage peeled (70 – 85%) while those grown at the Harris ranch had the lowest percentage peeled (20 to 30%). Although these locational differences appear important, there was no significant correlation between peelability and soil characteristics or production system at a particular farm.

The effect of the microwave thermal treatment on ascorbic acid, lycopene and total phenolics, in particular, was determined prior to the initiation of this project on three samples of Halley 3155 cultivar tomatoes. The microwave process resulted in a decrease in the concentration of both ascorbic and dehydroascorbic acid, but did not significantly affect the concentration of lycopene or total phenolics. Average ascorbic acid and dehydroascorbic acid concentrations in the raw tomatoes were 0.408 and 0.161 mg/100g while in microwaved juice they were 0.333 and 0.102 mg/100g, respectively. Average total phenolics concentrations were 100.5 ug/g and 103.0 ug/g in the raw and microwaved juices, respectively. Although a reduction in ascorbic acid resulted from the microwave process, it was felt that this preparative step was quite desirable for inactivation of enzymes that might cause an even greater reduction in quality, therefore it was utilized.

Results for the physico-chemical properties of conventionally and organically grown tomatoes are presented by individual grower in Table 5. There were significant differences in moisture content between tomatoes produced under conventional or organic production systems at all farms with the exception of Rominger, however in two cases the moisture content of conventional tomatoes was higher and in one case the organic tomato moisture content was higher. Therefore there was no consistent trend determined.

Over all, °Brix (soluble solids) of the tomatoes was higher under organic production systems in the grower group studied (Table 5). However on inspection of LSD values between individual growers, it appears that °Brix was significantly higher under organic production systems at the Terranova and Rominger farms but not significantly different at the Harris and O'Neill farms. Although there were differences between growers, the interaction of production system x grower was not significant (Table 5).

Soluble solids content is an important tomato quality parameter, both from a sensory flavor standpoint and a tomato processing standpoint. Tomato flavor is generally determined by the content of soluble solids, acid (measured as titratable acidity and expressed as % citric acid) and presence of various flavor volatiles in the tomatoes. Some of these flavor volatiles are generated

1 through enzyme-catalyzed reactions (Anthon and Barrett 2003). Flavor volatiles were not analyzed
2 in this study. There are differing opinions on the relative importance of these factors which
3 contribute to flavor, but Jones and Scott (1984) found that flavor impact is related to total sugar and
4 acid.

5 Soluble solids are important not only in terms of their contribution to flavor, but also, as
6 mentioned above, in terms of their relationship to processing requirements. It is desirable to
7 produce tomatoes higher in soluble solids content because they will require less energy to evaporate
8 water to a target final °Brix content. Tomato juice (~4-6 °Brix) is typically concentrated to 27-31
9 °Brix prior to packaging for long-term bulk storage as tomato paste. The higher °Brix levels in
10 tomatoes produced under organic production systems by the four growers in this study will result in
11 reduced processing costs.

12 The individual ANOVA results (Table 5) indicated no significant differences in tomato pH
13 values between production system, but there were differences between growers. The pH values
14 were higher in organic tomatoes than conventional at two grower locations, Terranova and O'Neill,
15 but lower at the Rominger farm and the same at the Harris farm. Titratable acidity levels were
16 higher in organically grown tomatoes, which may combine with the higher °Brix levels to
17 contribute favorably to flavor impact (Jones and Scott 1984). As was the case with °Brix, in
18 comparing the LSD values for specific growers, tomato citric acid content was significantly higher
19 under organic production systems at the Terranova and Rominger farms, but not significantly
20 higher at the Harris or O'Neill farms. Therefore, there was a significant effect associated with
21 grower, but the production system x grower interaction was not.

22 The ratio of sugars to acids is something that may be used to indicate general flavor quality,
23 and the fact that organically grown tomatoes in this study had higher soluble solids and higher
24 titratable acidity indicates that they may have better flavor quality. The sensory evaluation study
25 carried out with consumer panelists, however, did not find a significant difference in the flavor
26 acceptability of the organic vs. conventionally grown tomatoes (data not shown). Such a
27 difference, if it exists, might be determined by a descriptive sensory panel. The use of such a panel
28 was beyond the scope of the present study, but may be something to consider in future research.
29 There was a significant difference in Bostwick consistency between production systems in the four
30 growers evaluated (Table 5), with organic production systems producing tomatoes with
31 significantly lower Bostwick values, indicating a higher level of viscosity. Looking at individual
32 growers, Bostwick values were significantly lower under organic production systems at the
33 Terranova and Rominger farms, but not significantly different from conventional tomatoes at the
34 other two farms. Bostwick is another important quality attribute in tomato processing, and it will
35 often dictate what type of product tomatoes are intended for. Generally, higher Bostwick values are
36 undesirable as this indicates a less viscous product. In some processed products, such as paste and
37 sauce, a low Bostwick value is desirable, and breeding programs often target these varieties of
38 tomatoes. Paste and sauce products are typically thick and do not separate on standing. For other
39 products, however, such as whole peeled tomatoes, diced tomatoes, tomato juice and tomato soup,
40 having a viscous product with a low Bostwick value is not as critical. With these products, the
41 target quality parameters are flavor and color, and the viscosity of the product is not so important.
42 Therefore, from a Bostwick perspective, organically grown tomatoes may be more desirable than
43 conventionally grown tomatoes for paste and sauce products.

44 Catsup yield differed significantly by both grower and production system, but the
45 interaction of the two was not significant (Table 5). Within the four growers in this study, organic
46 production systems produced higher catsup yield than conventional production systems. Inspection
47 of the LSD values from the individual grower locations shows that catsup yield was significantly
48 higher under organic production systems at the Terranova and Rominger farms, but not different at
49 the Harris or O'Neill farms. Catsup yield is a value calculated from juice °Brix and juice Bostwick
50 values. High catsup yield is a desirable processing tomato quality attribute and indicates that

1 organically grown tomatoes may be better suited for producing catsup than conventionally grown
2 tomatoes.

3 4 Grower Specific Differences in Color of Conventionally and Organically Grown Tomatoes

5 Tomato color was determined using LED and Hunter tristimulus L, a, and b measurements
6 and USDA tomato paste color value was calculated. There was a significant difference at the $p <$
7 $.05$ level in LED values for tomato color both between the two production systems and the four
8 growers (Table 6). Inspection of LSD values of the LED color scores from individual growers
9 revealed that tomatoes were significantly more red when grown under conventional production
10 systems at the Terranova and Harris farms but not significantly different from organically grown
11 tomatoes at the Rominger and O'Neill farms. In the California processing tomato industry, loads of
12 harvested tomatoes are inspected by the Processing Tomato Advisory Board, a third party
13 inspection agency, prior to processing. A determination of whether the load is acceptable or not,
14 and the price the grower is paid, are dictated by the grade given.

15 A more robust measurement of color is accomplished through use of tristimulus
16 colorimeters. The a value is typically the most indicative for the intensity of red color in tomatoes,
17 with higher a values more desirable. Table 6 indicates there were significant differences between
18 growers ($p < .001$) in L and a value, and in the calculated USDA color score. The b value was
19 significantly different between growers at the $p < .05$ level. There were significant differences
20 between production system at the $p < .05$ level for both b value and USDA color score. Inspection
21 of the tomato color differences between individual growers shows that that there were no significant
22 differences in L value, but conventionally grown O'Neill tomatoes and organically grown
23 Rominger tomatoes had significantly higher a values, or more red color, than their counterparts.
24 The b values, which indicate increasing yellowness in the (+) direction, were significantly higher in
25 conventionally grown Terranova tomatoes but were not significantly different at the other three
26 farms. USDA color was significantly higher in organically grown Terranova tomatoes; but there
27 were no significant color differences between conventional or organic fruit grown at the other three
28 farms.

29 It may be most important to evaluate the results in terms of either LED value (currently used
30 by the California tomato industry at inspection stations) or the a value (most common analytical
31 laboratory parameter). In so doing, the LED result from the individual ANOVA statistics would
32 conclude that conventionally grown fruit are more red, while the a value would show mixed results
33 due to a significant interaction of production system and grower.

34 35 Grower Specific Differences in Nutrient Content of Conventionally and Organically Grown 36 Tomatoes

37 Nutrient levels were also associated with both the production system (organic vs.
38 conventional) and the individual grower (Table 7). Total ascorbic acid content (reduced ascorbic
39 acid or AA and dehydroascorbic acid or DHA) was generally lower in microwaved tomato juice as
40 compared to raw. In almost all comparisons, the sum of reduced and oxidized forms of ascorbic
41 acid was higher in tomatoes grown on conventional fields than those grown on organic at the same
42 grower location.

43 On two farms, Terranova and Harris, conventional production systems produced higher
44 values of ascorbic acid and lycopene in the microwaved juice. Terranova organically grown
45 tomatoes had significantly lower levels of dehydroascorbic acid in the microwaved juice.
46 Organically grown O'Neill tomatoes had significantly higher levels of lycopene, which is in
47 contradiction to the lycopene results from the Terranova and Harris farms. Lycopene may be one
48 attribute that shows an interaction between production system and grower. Total phenolics levels in
49 the microwaved juice samples were significantly higher in organically grown tomatoes from one

1 grower, O'Neill, but on two other farms (Terranova and Harris), the conventionally grown fruit had
2 significantly higher levels of total phenolics.

3 Interestingly, tomatoes grown at the Rominger farm consistently showed no difference in
4 lycopene, ascorbic acid or total phenolics in the microwaved juice as an effect of production
5 system. On the other hand, these three nutrients were always higher in conventional tomatoes
6 grown at Terranova and Harris, and conversely higher (or the same) in organic tomatoes grown at
7 O'Neill farm. If one examines the soil evaluations (Table 1), the Terranova and Harris Farms both
8 had sandy loam soil, while the O'Neill tomatoes were grown on clay. The Rominger tomatoes
9 (which showed no difference in nutrients) had a combination of silty clay and loam. There were
10 also some differences in tomato cultivar and plant type (Table 2), and in fertilizers and pesticides
11 used (Table 3) at these farms. The effect of organic and conventional production systems on
12 nutrient levels requires additional investigation because the results appear to have an interaction
13 with the specific farm studied.

14 15 Characterization of Soil Properties in Organic and Conventional Production Systems

16 The soil analysis was based on the four field locations where there were matched soil and
17 quality samples taken; and it was not possible to pool data for "all organic" or "all conventional".
18 Nor was it statistically valid to pool data for growers to determine the effects of these production-
19 related factors.

20 Three principal components described the soil characteristics of the conventional and
21 organic farms that were studied. These soil samples were taken in each field after fruit set and prior
22 to planting. There was a significant difference between the Terranova farm and others in an
23 ANOVA with PC1 as the outcome. Sodium content was relatively high in both conventional and
24 organic soils at this farm, and Mg was high in organic soils, as compared to the other three farms.
25 PC2 separated the four different growers, with the Harris farm being lower than others in some soil
26 characteristics. PC3 separated the organic and conventional production systems.

27 Soils from conventional tomato production fields were generally higher in pH, exchange
28 capacity (EC), Ca, Na and Cl. Organic production soils were higher in HCO₃, P, Mn, Fe and
29 moderately higher in K, N and Cu. Therefore, it was possible to distinguish the four organic fields
30 from their conventional counterparts in terms of these specific soil parameters. In a study
31 comparing the major and trace elements in organic and conventional Danish agricultural crops,
32 there were significant differences in element concentration mean values between organically and
33 conventionally grown onions and peas (Gundersen and others 2000). These investigators also used
34 PCA to separate the elements measured into two groups according to the cultivation method. In the
35 case of onions, there were differences in some of the same elements evaluated in this study, in
36 particular Ca (calcium). Svec and others (1976) found organically grown tomatoes to be higher in K
37 than conventionally grown. Further research on the effects of these specific components on tomato
38 quality and nutritional value may allow for an understanding of the mechanism by which organic
39 production systems affect these attributes.

40 The ANCOVA (Analysis of Covariance) model using the PCs was compared to an ANOVA
41 statistical model, in which all 22 soil characteristics were utilized. In 10 out of 18 outcomes
42 (quality attributes), the models were not significantly different, according to the lack-of-fit test,
43 meaning the 3 PCs were "representative enough" of the farms for those outcomes. These 10 quality
44 attributes included: lycopene, dehydroascorbic acid in the microwaved juice, catsup yield, Brix,
45 pH, Bostwick, LED, a value, b value, and USDA value. The 8 attributes where it was not possible
46 substitute the PCs for model with all of the soil data were: T.A., ascorbic acid, dehydroascorbic
47 acid, L value, raw and microwaved juice polyphenols. This means that it was possible to represent
48 the differences in organic and conventional production systems by PC3, in particular, for some
49 quality parameters, but not others.

1
2 **Conclusions**

3 The goal of this study was to compare the quality and nutritional value of processing
4 tomatoes grown on matched organic and conventional farms at four commercial grower locations.
5 While we were successful in our evaluation of tomatoes grown under these “real life” conditions,
6 because we were only able to include four growers in the study, our conclusions are restricted to
7 this specific group of growers. It was not possible to make statements concerning the differences
8 between quality of organic and conventionally grown tomatoes. In retrospect, our statistical
9 analysis would have been stronger had we selected a larger number (6-10) of growers and a smaller
10 number (4) of sampling locations per field.

11 Within the grower group that we worked with, there were differences even within
12 production system (e.g. organic or conventional) in fertilizer and irrigation systems, methods of pest
13 control and tomato cultivar planted. However, these preliminary results on a limited number of
14 commercial growers indicated that there may be potential advantages to the use of organically
15 grown tomatoes because of higher levels of soluble solids, titratable acidity and consistency. This
16 study of four growers indicated that both higher flavor impact and reduced energy requirement for
17 concentration of juice may result from use of organically grown tomatoes. Our study showed that
18 conventionally produced tomatoes, on the other hand, were more red in color and the microwaved
19 juice was higher in ascorbic acid and total phenolics.

20 One important result of this work was the finding that the agricultural production system is a
21 critical factor in determining the quality of fruit produced. The potential exists to affect both
22 positive and negative attributes of fruit quality through a better understanding of the production
23 system, whether it be conventional, organic or something else. Through this type of study it may be
24 possible to identify major factors that influence quality.

25 In the future we will attempt to collect data from a larger number of commercial growers in
26 order to improve our ability to make statistically valid conclusions about global differences between
27 organic and conventionally grown tomatoes. In addition, we would like to attempt to correlate our
28 findings in commercial systems with those in university-based research plots, where the production
29 system may be better controlled in terms of fertilizer applications, soil type, irrigation system and
30 methods of pest control. Only when we are able to separate out the influence of grower-specific
31 inputs will it be possible to conclusively state whether there are differences in nutritional quality
32 between organic and conventional systems or not.

References

- Anthon A, Barrett, DM. 2003. Thermal inactivation of lipoxygenase and hydroperoxytrieneoic lyase in tomatoes. *Food Chem* 81, 2:275-279.
- AOAC International. 2000. Official methods of analysis. 17th edition ed. Gaithersburg, MD: AOAC Int.
- Asami DK, Hong YH, Barrett DM, Mitchell AE. 2003. A comparison of the total phenolic and ascorbic acid contents of freeze-dried and air-dried Marionberry, strawberry and corn grown using conventional, organic and sustainable agricultural practices. *JAgFdChem* 51:1237-41.
- Barrett DM. 2001. Tomato attributes and their correlation to peelability and product yield. *Acta Horticulturae* 542:65-74.
- Barrett DM, Anthon GE. 2001. Lycopene content of California-grown tomato varieties. *Acta Horticulturae* 542:165-74.
- Bourn D, Prescott J. 2002. A comparison of the nutritional value, sensory qualities and food safety of organically and conventionally produced foods. *Crit Rev Food Sci Nutr* 42:1-34.
- Caris-Veyrat C, Amiot MJ, Tyssandier V, Grasselly D, Buret M, Mikolajczak M, Guillard JC, Bouteloup-Demange C, Borel P. 2004. Influence of organic versus conventional agricultural practice on the antioxidant microconstituent content of tomatoes and derived purees; consequences on antioxidant plasma status in humans. *J Agric Food Chem* 52:6503-9.
- Chassy, A., Bui, L., Renaud, E.N.C., Van Horn, M., Mitchell, A. 2006. Three-Year Comparison of the Content of Antioxidant Microconstituents and Several Quality Characteristics in Organic and Conventionally Managed Tomatoes and Bell Peppers. *J Agric Food Chem* 54 (21), 8244 - 8252.
- Diver S. 2000. Nutritional Quality of Organically Grown Food, ATTRA (Appropriate Technology Transfer for Rural Areas). Arkansas, U.S.
- Economic Research Service. 2005a. Data: Organic production, 1992-2003.
- Economic Research Service. 2005b. Briefing room: Organic farming and marketing. United States Department of Agriculture.
- Garcia E, Barrett DM. 2006. Evaluation of processing tomatoes from two consecutive growing seasons: Quality attributes, peelability and yield. *J Food Proc Preserv.* 30:20-36.
- Gunderson V, Bechmann IE, Behrens A, Sturup, S. 2000. Comparative investigation of concentrations of major and trace elements in organic and conventional Danish agricultural crops. 1. Onions (*Allium cepa* Hysam) and peas (*Pisum sativum* Ping Pong). *J Agric Food Chem* 48:6094-6102.
- Heaton S. 2001. Organic Farming, Food Quality and Human Health: A Review of the Evidence Bristol, U.K: Soil Association.
- Johnson, R. A., Wichern, D. W. 1998. *Applied Multivariate Statistical Analysis* (4th ed.), Upper Saddle River, NJ: Prentice Hall.
- Jones RA, Scott SJ. 1984. Genetic potential to improve tomato flavor in F[1] hybrids. *J Am Sci Hort Sci.* 109, 3:318-321.
- Klonsky K, Richter K. A statistical picture of California's organic agriculture: 1998-2003. Agricultural Issues Center, UC Davis.
- Kutner M.H., Nachtsheim C.J., Neter J., Li W. 2004. *Applied Linear Statistical Models* (5th ed.), Homewood, IL: Irwin.
- Latapi L, Barrett DM. 2006. Influence of pre-drying treatments on quality and safety of sun-dried tomatoes. Part II. Effects of storage on nutritional and sensory quality of sun-dried tomatoes pretreated with sulfur, sodium metabisulfite or salt. *J Food Science* 71(1):32-7.
- Leonard S, Buhlert J, Marsh G, Wolcott T, Heil JR. 1980. Procedures for evaluating utilization potential and quality in processing tomatoes and tomato products. Dept. Food Science and Technology, U.C. Davis, Davis, CA. 72 p.
- Microsoft. 2000. Excel for Windows computer software.

- Organic Trade Association. 2004. OTA's 2004 Manufacturer Survey.
- Pollard J, Kirk SFL, Cade JE. 2002. Factors affecting food choice in relation to fruit and vegetable intake: A review. *Nutrition Research Reviews* 15:373-87.
- Reganold JP, Glover JD, Andrews PK, Hinman HR. 2001. Sustainability of three apple production systems. *Nature* 410(April).
- Svec, LV, Thoroughgood, CA, Mok, HCS. 1976. Chemical evaluation of vegetables grown with conventional or organic soil amendments. *Commun Soil Sci Plant Anal* 7:213-28.
- The Soil Association. 1999. The organic food and farming report. Bristol, UK: The Soil Association.
- Valero, C., Cristoso, C.H., Garner, D., Bowerman, E., Slaughter, D. 2003. Introducing nondestructive flesh color and firmness sensors to the tree fruit industry. *Acta Hort.* 604, 597-600.
- Veberic R, Trobec M, Herbinger K, Hofer M, Grill D, Stampar F. 2005. Phenolic compounds in some apple (*Malus domestica* Borkh) cultivars of organic and integrated production. *J Sci Food Agric* 85:1687-94.
- Weibel F, Bickel R, Leuthold S, Alfoldi T. 2000. Are organically grown apples tastier and healthier? A comparative field study using conventional and alternative methods to measure fruit quality. *Acta Horticulturae* 57:417-26.
- Winter CK. 2006. Organic foods. *J. Food Sci* 10:44-48.
- Woese K, Lange D, Boess C, Bogel KW. 1997. A comparison of organically and conventionally grown foods - results of a review of the relevant literature. *J Sci Food Agric* 74:281-93.
- Worthington V. 1998. Effect of agricultural methods on nutritional quality: a comparison of organic with conventional crops. *Alternative Therapies Health Med* 4(1):58-69.
- Young, JE, Zhao X, Carey EE, Welti R, Yang S-S, Wang W. 2005. Phytochemical phenolics in organically grown vegetables. *Mol Nutr Food Res* 49:1136-42.

Acknowledgements

The authors gratefully acknowledge the contributions of D. Cameron, Terranova Ranch, Inc., L. Chrisco, Harris Farms, Inc., J. Rominger, D.A Rominger & Sons, Inc., C. Stubblefield, O'Neill Farming Co. Inc., USDA-NRCS Soil Scientists Kerry Arroues and Jim Komar, and research consultants Dr. John Reganold, Professor of Soil Science, Washington State University, and Dr. Frank Zalom, Professor of Entomology, University of California at Davis. We would like to thank Small Planet Foods for their support of this project.

Table 1. USDA-NRCS evaluation of soil properties in organic and conventional fields at four farms in California.

Grower	Location	Production System	Soil Type	Horizon	Depth (centimeters)	Soil Texture				
Harris Farms	Coalinga	Organic	Kimberlina Sandy Loam	Ap	0 - 17.8	sandy loam				
				A	17.8 - 50.8	sandy loam				
				C1	50.8 - 63.5	sandy loam				
				C2	63.5 - 76.2	sandy loam				
				C3	76.2 - 88.9	sandy loam				
				C4	88.9 - 101.6	sandy loam				
				Conventional	Kimberlina Sandy Loam	Ap	0 - 17.8	sandy loam		
		A	17.8 - 50.8			sandy loam				
		C1	50.8 - 71.1			silt loam				
		C2	71.1 - 101.6			fine sandy loam				
		O'Neill Farming Co.	Five Points			Organic	Ciervo Clay	Ap	0 - 38.1	clay
								Bw	38.1 - 58.4	clay
								Bk	58.4 - 83.8	clay loam
				Bzn	83.8 - 101.6			clay loam		
Conventional	Ciervo Clay			Ap	0 - 30.5			clay loam		
				Bw	30.5 - 45.7			clay		
				Bk1	45.7 - 68.6			clay		
				Bk2	68.6 - 81.3	clay loam				
				Bk3	81.3 - 101.6	clay loam				
				Terranova Ranch	Helm	Organic	Pond Fine Sandy Loam	Ap1	0 - 27.9	loam
								Ap2	27.9 - 50.8	loam
Btkn1	50.8 - 76.2							clay loam		
Btkn2	76.2 - 91.4							clay loam		
Bk	91.4 - 101.6							loam		
Conventional	Pond Fine Sandy Loam	Ap1	0 - 27.9					loam		
		Ap2	27.9 - 45.7					loam		
		Btkn	45.7 - 71.1			clay loam				
		Bk1	71.1 - 86.4			loam				
		Bk2	86.4 - 101.6			loam				
		D.A. Rominger & Sons	Winters			Organic	Brentwood Silty Clay Loam	Ap1	0 - 17.8	silty clay loam
								Ap2	17.8 - 30.5	silty clay loam
								heavy silty clay		
B21	30.5 - 45.7							loam		
B22	45.7 - 55.9			silty clay loam						
B3	55.9 - 81.3			silt loam						
BC	81.3 - 106.7			silty clay loam						
				heavy silty clay						
C	106.7 +			loam						
Conventional	Brentwood Silty Clay Loam			Ap	0 - 27.9			silty clay loam		

B21	27.9 - 50.8	silty clay loam
B22	50.8 - 61	silty clay loam
B3	61 - 96.5	heavy silt loam
C1	96.5 - 111.8	silty clay loam/clay loam
C2	111.8 +	heavy clay loam

¹Soil evaluation carried out by USDA-NRCS Soil Scientists Kerry Arroues and Jim Komar

Table 2. Organic and conventional tomato production system characteristics for four farms in California.

Grower and Production System	Location	Tomato Cultivar	Planting Type	Planting Date	Irrigation Method	Organic Certification	Years Farmed	Harvest Sample Date	% Ripe	Sensory Sample Date	Yield (kg/ha)
									Fruit by Weight		
Harris Farms	Coalinga										
Organic		HyPeel 45	Transplants	4/4/03	Drip	1996	30+	8/5/2203	95	8/5/2003	68.38
Conventional		HyPeel 45	Transplants	4/10/03	Drip	-	30+	8/5/2203	90	8/5/2003	59.64
O'Neill Farming Co.	Five Points										
Organic		Rogers 1570	Seed	1/30/03 - 1/31/03	Furrow	1991	50+	7/23/2003	87	7/22/2003	85.42
Conventional		Rogers 1570	Seed	1/31/03 - 2/4/03	Furrow	-	50+	7/22/2003	96	7/22/2003	108.51
Terranova Ranch	Helm										
Organic		Bos 315	Transplants	3/22/2003	Furrow	1997	45+	7/29/2003	89	7/29/2003	77.80
Conventional		Bos 315	Transplants	3/22/2003	Furrow	-	45+	7/22/2003	87	7/29/2003	76.45
D.A. Rominger & Sons	Winters										
Organic		HM 830	Transplants	3/31/03 - 4/1/03	Furrow	1996	70+	7/30/2003	94	8/6/2003	96.41
Conventional		HM 830	Transplants	3/31/03	Furrow	-	70+	6/8/2003	96	8/6/2003	109.86

Table 3. Pesticide and fertilizer use by grower and production system.

Grower	Production System	Soil Fertility Materials/Methods	Fertility Materials Rate per Hectare	Pest Control Materials/Methods	Pest Control Materials Rate per Hectare (a.i.)	
Harris Farms	Organic	cow manure	13,452 kg/ha	copper hydroxide	1.73 kg/ha	
		chicken manure	11,210 kg/ha	<i>Bacillus thurengiensis</i>	0.064 kg./ha	
		compost	6,726 kg/ha	sulfur	33.6 kg/ha	
	Conventional	urea ammonium nitrate	19.1 kg/ha (N)	rimsulfuron herbicide	.0058 kg/ha	
		urea ammonium nitrate	168 kg/ha (N)	rimsulfuron herbicide	.0058 kg/ha	
		10-34-0 (NPK)	1.26 kg/ha (N); 4.8 kg/ha (P ₂ O ₅)	<i>Bacillus thurengiensis</i>	.01 kg/ha	
			tebufenozide insecticide	.175 kg/ha		
			<i>Bacillus thurengiensis</i>	.064 kg/ha		
			indoxacarb insecticide	.074 kg/ha		
			sulfur	38.5 kg/ha		
indoxacarb insecticide	.074 kg/ha					
pyraclostrobin fungicide	.168 kg/ha					
esfenvalerate insecticide	.055 kg/ha					
O'Neill Farming Co.	Organic	winter wheat cover crop		<i>Bacillus thurengiensis</i>	.023 kg/ha	
		Compost	11,210 kg/ha			
		Bird Guano	41.5 kg/ha			
		Chilean Nitrate	24.7 kg/ha			
	Conventional	11-52-0 (NPK)	37 kg/ha (N); 175 kg/ha (P ₂ O ₅)	rimsulfuron herbicide	.0106 kg.ha	
			aqua ammonia	84 kg/ha (N)	trifluralin herbicide	.713 kg/ha
		10-34-0 (NPK)	1.4 kg/ha (N); 4.8 kg/ha (P ₂ O ₅)	copper hydroxide	1.34 kg/ha	
			urea ammonium nitrate	123.3 kg/ha (N)	maneb fungicide	1.35 kg/ha
			calcium ammonium nitrate	23.5 kg/ha (N)	sulfur	43.9 kg/ha
			urea ammonium nitrate	97.5 kg/ha (N)	indoxacarb insecticide	.074 kg/ha
Terranova Ranch	Organic	chicken manure	15,694 kg/ha	copper hydroxide	1.73 kg/ha	
		chicken compost	38,114 kg/ha	sulfur	43.7 kg/ha	
	Conventional	11-52-0 (NPK)	31.4 kg/ha (N); 145.7 kg/ha (P ₂ O ₅)	rimsulfuron herbicide	.058 kg/ha	
			Mg & B micronutrients	copper hydroxide	1.37 kg/ha	
		urea ammonium nitrate	140 kg/ha (N)	sulfur	43.7 kg/ha	
urea ammonium nitrate	134.5 kg/ha (N)					
urea ammonium nitrate	67.3 kg/ha lbs./A (N)					

D.A. Rominger & Sons	Organic	turkey manure	15,694 kg/ha	copper hydroxide	3.45 kg/ha
				copper hydroxide	3.45 kg/ha
				sulfur	33.6 kg/ha
	Conventional	11-52-0 (NPK)	12.3 kg/ha (N); 58.3 kg/ha (P ₂ O ₅)	glyphosate herbicide	.91 kg/ha
				8-24-5 (NPK)	1.1 kg/ha (N); 3.3 kg/ha (P ₂ O ₅)
		zinc micronutrient		maneb fungicide	.598 kg/ha
		28-0-0-5 (NPKS)	15.7 kg/ha (N)	copper hydroxide	1.75 kg/ha
		urea ammonium nitrate	78.5 kg/ha (N)	esfenvalerate insecticide	.185 kg/ha
				rimsulfuron herbicide	.058 kg/ha
				maneb fungicide	.598 kg/ha
		copper hydroxide	1.75 kg/ha		
		sulfur	33.3 kg/ha		

Table 4. Multivariate analysis of variance (MANOVA) F values for grower, production system and interaction of grower and production system

	Wilk's Lamda	Pillai's Trace	Hotelling Lawley Trace	Roy's Greatest Root
Grower	< .001	< .001	< .001	< .001
Production System	< .001	< .001	< .001	< .001
Grower x Production System	< .001	< .001	< .001	< .001

Table 5. Physico-chemical properties of conventionally and organically grown tomatoes

Grower	Moisture		°Brix		pH		T.A.		Bostwick		Catsup		Texture	
	(%)						(%)		(cm)		Yield (kg)		(N) ¹	
Harris Farms														
Conventional	94.57	a ²	4.86	de	4.65	a	0.26	c	16.70	ab	370.80	cd	188.4	d
Organic	93.05	cd	5.11	cd	4.64	a	0.27	c	16.18	b	385.28	bc	155.7	d
O' Neill Farming Co,														
Conventional	93.28	c	5.06	d	4.47	c	0.31	b	16.13	b	383.89	bc	202.7	c
Organic	94.76	a	5.03	d	4.55	b	0.30	b	15.68	b	389.95	bc	200.0	c
Terranova Ranch														
Conventional	94.30	ab	5.56	b	4.32	d	0.31	b	16.30	b	399.49	b	238.1	c
Organic	92.84	d	5.96	a	4.52	bc	0.37	a	14.13	c	438.36	a	185.4	d
D.A. Rominger & Sons														
Conventional	94.48	a	4.66	e	4.70	a	0.21	d	18.22	a	344.73	d	412.8	a
Organic	94.35	ab	5.39	bc	4.52	bc	0.25	c	15.05	bc	409.06	b	300.3	b
Pr > F														
Grower			< .001		< .001		< .001		0.0539		< .001			
Production system			< .001		< 0.1553		< .001		0.0005		< .001		0.12	
Production System X			< .007		< .001		0.0029		0.0844		0.0146			
Grower														

¹Texture is reported in Newtons, which is the amount of force required to accelerate a body with a mass of one kilogram at a rate of one meter per second squared. 1 N = 1 kg m/s²

²Means in a column followed by the same letter are not significantly different at the 5% level of probability.

Table 6. Color of conventionally and organically grown tomatoes.

Grower	LED		L		Hunter a		b		USDA Color	
Harris Farms										
Conventional	22.3	d ¹	23.14	bc	26.12	bc	15.00	ab	44.4	d
Organic	21.6	ab	23.13	bc	26.35	bc	15.08	ab	44.5	d
O' Neill Farming Co.										
Conventional	21.7	bc	22.36	d	27.53	a	15.08	bc	45.6	b
Organic	21.8	abc	23.09	cd	26.24	bc	14.64	c	45.3	bc
Terranova Ranch										
Conventional	22.1	d	23.17	bc	27.77	a	15.24	a	45.6	b
Organic	21.3	a	23.42	ab	27.26	a	14.40	d	46.7	a
D.A. Rominger & Sons										
Conventional	22.5	d	23.61	a	25.87	c	14.67	c	44.8	cd
Organic	22.1	cd	23.51	a	26.63	b	14.84	bc	45.3	bc
Pr > F										
Grower	0.0029		< .001		< .001		0.0030		< .001	
Production system	0.0003		0.1813		0.2786		0.0020		0.0170	
Production System X	0.102		0.4024		< .001		< .001		0.0047	
Grower										

¹Means in a column followed by the same letter are not significantly different at the 5% level of probability.

Table 7. Nutrient concentration (dry weight) of conventionally and organically grown tomatoes

Grower	Ascorbic Acid		Dehydro		Total	
	raw ¹ ug/g		Ascorbic Acid raw ug/g		AA+DHA raw ug/g	Ascorbic Acid microwaved ug/g
Harris Farms						
Conventional	1187	ab ²	2306	b	3493	2211
Organic	1133	ab	1928	c	3061	1810
O'Neill Farming Co.						
Conventional	1739	d	1756	cd	3495	1155
Organic	158	d	2448	b	2606	1302
Terranova Ranch						
Conventional	863	bc	1918	c	2781	2016
Organic	638	c	1587	d	2225	1319
D.A. Rominger & Sons						
Conventional	1372	a	3031	a	4403	2455
Organic	1153	ab	2242	b	3395	2258
Pr > F						
Grower	< .001		< .001			< .001
Production system	0.15		0.0125			< .001

¹Nutrient concentration was determined in raw and microwaved tomato juice.

²Means in a column followed by the same letter are not significantly different at the 5% level of probability.

